

FORUM

The LEDA Traitbase: a database of life-history traits of the Northwest European flora

M. Kleyer^{1*}, R.M. Bekker^{1,2}, I.C. Knevel^{1,2}, J.P. Bakker², K. Thompson³, M. Sonnenschein^{4,5}, P. Poschlod⁶, J.M. van Groenendael⁷, L. Klimeš⁸, J. Klimešová⁸, S. Klotz⁹, G.M. Rusch¹⁰, M. Hermy¹¹, D. Adriaens¹¹, G. Boedeltje⁷, B. Bossuyt¹⁸, A. Dannemann¹, P. Endels¹¹, L. Götzenberger⁹, J.G. Hodgson¹³, A-K. Jackel⁶, I. Kühn⁹, D. Kunzmann¹, W.A. Ozinga^{7,12}, C. Römermann⁶, M. Stadler^{4,5}, J. Schlegelmilch⁵, H.J. Steendam², O. Tackenberg⁶, B. Wilmann¹⁰, J.H.C. Cornelissen¹⁴, O. Eriksson¹⁵, E. Garnier¹⁶ and B. Peco¹⁷

¹Carl von Ossietzky University of Oldenburg, Landscape Ecology Group, P.O. Box 2503, 26111 Oldenburg, Germany;

²University of Groningen, Community and Conservation Ecology Group, P.O. Box 14, 9750 AA Haren, The Netherlands;

³Sheffield University, Department of Animal and Plant Sciences, Sheffield S10 2TN, United Kingdom; ⁴Carl von Ossietzky University of Oldenburg, Department of Computer Science, Uhlhornsweg D-26111 Oldenburg, Germany; ⁵OFFIS e.V.,

Escherweg 2, 26121, Oldenburg, Germany; ⁶University of Regensburg, Institute of Botany, D-93040 Regensburg, Germany; ⁷Radboud University, Aquatic Ecology & Environmental Biology, Toernooiveld 1, 6525 ED Nijmegen, The

Netherlands; ⁸Institute of Botany (ASCR), Section of Plant Ecology, Dukelská 135, 379 82 Třebon, Czech Republic;

⁹Helmholtz-Centre for Environmental Research – UFZ, Department of Community Ecology (BZF), Theodor-Lieser-Strasse 4, D-06120 Halle (Saale), Germany; ¹⁰Norwegian Institute for Nature Research (NINA), 7485 Trondheim, Norway;

¹¹University of Leuven, Division Forest, Nature & Landscape Research, Celestijnenlaan 200 E, 3001 Leuven, Belgium;

¹²Alterra Wageningen UR, P.O. Box 47, 6700 AA Wageningen, The Netherlands; ¹³Peak Science and Environment, Station House, Hathersage, S32 1BA, Derbyshire, United Kingdom; ¹⁴Institute of Ecological Science, Department of Systems

Ecology, Vrije Universiteit Amsterdam, De Boelelaan 1087, 1081 HV Amsterdam, The Netherlands; ¹⁵Department of Botany, Stockholm University, 106 91 Stockholm, Sweden; ¹⁶Centre d'Ecologie Fonctionnelle et Evolutive (UMR5175),

CNRS, 1919 route de Mende, 34293 Montpellier Cedex 5, France; ¹⁷Universidad Autónoma de Madrid, Ecology Department, 28049 Cantoblanco, Madrid, Spain; and ¹⁸Ghent University, Terrestrial Ecology Unit, K. L. Ledeganckstraat 35 B-9000 Ghent, Belgium

Summary

1. An international group of scientists has built an open internet data base of life-history traits of the Northwest European flora (the LEDA-Traitbase) that can be used as a data source for fundamental research on plant biodiversity and coexistence, macro-ecological patterns and plant functional responses.

2. The species-trait matrix comprises referenced information under the control of an editorial board, for ca. 3000 species of the Northwest European flora, combining existing information and additional measurements. The data base currently contains data on 26 plant traits that describe three key features of plant dynamics: persistence, regeneration and dispersal. The LEDA-Traitbase is freely available at www.leda-traitbase.org.

3. We present the structure of the data base and an overview of the trait information available.

4. *Synthesis.* The LEDA Traitbase is useful for large-scale analyses of functional responses of communities to environmental change, effects of community trait composition on ecosystem properties and patterns of rarity and invasiveness, as well as linkages between traits as expressions of fundamental trade-offs in plants.

Key-words: age of first flowering, buoyancy, clonal traits, canopy height, dispersal, functional ecology, plant life span, plant functional traits, seed weight, seed number, SLA, soil seed bank

*Correspondence author. E-mail: michael.kleyer@uni-oldenburg.de

Introduction

The immense variation in plant form and life history has always intrigued botanists, plant geographers and ecologists. From the middle of the 19th century, interest in disentangling relationships between plant biological traits and the environment has steadily developed, resulting in a wealth of descriptions of plant morphology as adaptations to climate and soil factors (Du Rietz 1931). This interest evolved into compilations of biological knowledge for individual plant species (e.g. Kirchner *et al.* 1908–1936; the Biological Flora of the British Isles series published in this journal; Rabotnov 1974–1990 for Russia). A further step has been taken more recently, to build up digital data bases to synthesise information on plant traits. For instance, the GLOPNET data base (Wright *et al.* 2004) covers chemical, structural and physiological traits of leaves for a large number of species worldwide. Seed weight data are now available for > 10⁴ species (Flynn *et al.* 2006), and other data bases offer bibliographic data for selected communities (e.g. APIRS for aquatic plants (<http://plants.ifas.ufl.edu/>)) or provide taxonomic information together with some selected traits (e.g. USDA 2006).

Within Europe, knowledge of traits for individual species is growing fast, but information remains scattered over many sources, including dozens of different journals, large monographs and floras. The sources are available in various languages and distributed across many countries, collected and stored in different ways, and are not mutually integrated. Standardisation of trait definitions and measurements is often poor among species and studies. Trait data can also be retrieved from various data bases. However, currently accessible data bases are often restricted to certain regions, and cover only a limited number of species or traits.

A trans-national initiative has therefore aimed at designing and filling a species-trait matrix for the NW European flora that would be freely retrievable on the Worldwide Web (Knevel *et al.* 2003). The LEDA Traitbase (www.leda-traitbase.org), which uses a European consolidated species list, is concerned with pooling existing data bases, compiling new information from published data and closing knowledge gaps through extensive new measurements across several NW European countries. It consists of a relational data base linking species with traits and reference information about data source, location, habitat and trait measurement protocol on three core sets of traits: (i) persistence (vegetative) traits such as leaf, stem and clonal growth characteristics; (ii) regeneration traits such as seed production, seed longevity and (iii) dispersal traits such as seed weight, dispersal vectors, floating capacity and vertical terminal velocity of propagules.

The general objectives of the LEDA project were announced in Knevel *et al.* (2003). The present article describes the scope and architecture of the data base, the methods of collecting data and the plant life-history traits that are covered by the LEDA Traitbase. Additionally, a brief overview of applications illustrates the value of trait data bases in general, and the LEDA Traitbase in particular, for research in functional ecology.

Framework of the LEDA Traitbase

TRAITS IN THE LEDA TRAITBASE

Traits covered by the LEDA Traitbase were selected according to two major criteria: (1) relevance for persistence, regeneration and dispersal as key functions for survival in patterned landscapes and (2) trait data available for the flora of Northwest Europe, either in published sources or in unpublished data bases maintained by the project partners. As LEDA was designed as a compilation of data for a large number of species, we had to exclude traits for which only a small number of records for the Northwest European flora could be expected (e.g. relative growth rate or leaf life span). Table 1 shows an overview of the traits in the LEDA Traitbase together with associated functions and selected references, whereas Table 2 describes the categories or units of measurement, and actual number of species and records for the traits (version 1, 2007). More detailed information on the trait definitions is available in the Appendix S1 in Supporting Information.

Many trait data now available in LEDA for Northwest Europe had already attracted considerable attention in functional ecology (e.g. canopy height, seed number, seed mass, see Table 1) and are, at least in part, available elsewhere (e.g. Ellenberg *et al.* 1991 for life form; Flynn *et al.* 2006 for seed mass). For other traits, the LEDA Traitbase may be a unique source of data. For instance, seed bank longevity of many species was poorly known previously. The LEDA Project has improved this knowledge quite substantially from 21 071 records on 1189 species in the data base of Thompson *et al.* (1997) to 44 353 records covering 1787 species in total in the LEDA Traitbase (Table 2).

The LEDA Traitbase also includes data on clonal growth and dispersal traits that are rarely available elsewhere. The morphological traits characterising clonal growth serve as indicators for vegetative multiplication, persistence and vegetative regeneration subsequent to damage (Klimeš *et al.* 1997; Klimešová & Martínková 2004). The data available in the LEDA Traitbase that are related to clonal growth encompass a categorisation of clonal growth organs, bud bank vertical distribution and seasonality (Klimešová & Klimeš 2007), life span of a shoot, persistence of the connections between parent and offspring shoots, lateral spread and number of offspring shoots produced per year and per parent shoot. These traits indicate speed of lateral spread, rate of clonal multiplication and duration of possibility of mutual support inside interconnected parts of a clone.

Seed dispersal influences many key aspects of the biology of plants, but is inherently hard to measure (Cain *et al.* 2000). Since every species may be dispersed through different vectors and to different distances, we have measured traits related to dispersal potential (Poschlod *et al.* 2005). Terminal velocity is a relevant predictor for wind dispersal potential. If vertical air velocity exceeds 'terminal velocity' then the seed can be uplifted and dispersed for larger distances (Nathan *et al.* 2002; Tackenberg *et al.* 2003). Combined with terminal velocity,

Table 1. Overview of the traits in the LEDA Traitbase, their functional significance and related publications

The LEDA traits	Functional significance and related publications
<i>Persistence</i>	
Canopy height	Competitive ability (Westoby <i>et al.</i> 2002)
Leaf distribution along the stem, branching, shoot growth form	Competitive ability (Barkman 1988)
Leaf mass, leaf size, specific leaf area, leaf dry matter content	Growth rate, competitive ability, stress tolerance (Westoby <i>et al.</i> 2002)
Woodiness, stem specific density	Growth rate, investment in supporting structure (Ryser 1996)
Clonal growth organs, persistence of connection between parent and offspring shoots, number of offspring shoots per parent shoot per year, lateral spread	Competitive ability, persistence, clonal integration, storage (De Kroon & Van Groenendael 1997; Klimeš & Klimešová 2000; Vesik & Westoby 2004)
Bud bank – vertical distribution and seasonality	Response to disturbance (Bellingham & Sparrow 2000; Klimešová & Klimeš 2007)
<i>Regeneration</i>	
Plant growth form, plant life span, age of first flowering	Response to disturbance, establishment, invasiveness (Raunkiaer 1937; Rejmánek & Richardson 1996)
Seed number, seed shedding	Response to disturbance, establishment, dispersal (Leishman 2001; Bruun & Poschlod 2006)
Seed weight, size and shape	Dispersal, establishment (Grime <i>et al.</i> 1988; Westoby <i>et al.</i> 2002)
Seed bank longevity	Storage effects, response to disturbance (Bekker <i>et al.</i> 1998)
<i>Dispersability</i>	
Morphology of dispersal unit, seed releasing height	Wind dispersal, ecto- and endozoochorous dispersal (Van der Pijl 1972)
Dispersal vectors	Spectra of dispersal vectors for plants (Bonn <i>et al.</i> 2000)
Terminal velocity	Wind-dispersal (Tackenberg <i>et al.</i> 2003)
Attachment capacity of the dispersal unit, digestion survival	Ecto- and endozoochorous dispersal (Couvreur <i>et al.</i> 2004; Römermann <i>et al.</i> 2005)
Buoyancy	Dispersal in running water (Danvind & Nilsson 1997)

Table 2. Contents of the LEDA Traitbase, version 1

Trait name in data standards	Number of species	Number of records	Catalogue Number	Category or unit(s) of measurement
Plant growth form	2334	3154	1	Phanerophyte
			2	Chamaephyte
			3	Hemicryptophyte
			4	Cryptophyte
			4.1	Geophyte
			4.2	Helophyte
			4.2.1	Halophyte
			4.3	Hydrophyte
			5	Therophyte
			6	Liana
			7	hemi-epiphyte
Canopy height	2893	4934	8	Epiphyte
			9	vascular semi-parasite
Plant life span	2219	4293	10	vascular parasite
			11	mesophyte
				m
			1	summer annuals
			2	winter annuals
			3	strict monocarpic biennials and poly-annuals
			4	short-lived perennials (< 5 years)
			5	medium-lived perennials (5–50 years)
			6	long-lived perennials (> 50 years)
			7	perennials without any further detailed information
			Age of first flowering	1521
2	1 and 5 years			
3	> 5 years			
Leaf mass	1665	4472		mg
Specific leaf area (SLA)	2019	5941		mm ² mg ⁻¹
Leaf size	2054	5590		mm ²
Leaf dry matter content (LDMC)	1735	3451		mg g ⁻¹

Table 2. Continued

Trait name in data standards	Number of species	Number of records	Catalogue Number	Category or unit(s) of measurement
Woodiness & Stem specific density	3152	5300	1	woody
			1.1	hard wood
			1.2	soft wood
			2	semi-woody
			3	herbaceous (non-woody)
Shoot growth form	3118	5386		g cm ⁻³
			1	lianas, climbers and scramblers
			2	stem erect
			3	stem ascending to prostrate
			4	stem prostrate
			5	free-floating plants
			6	emergent, attached to the substrate
			7	floating leaves, attached to the substrate
Branching	2878	4055	8	submerged, attached to the substrate
			1	yes
			2	no
Leaf distribution along the stem	3491	5355	3	unknown
			1	rosette/tufted plant
			2	semi-rosette
			3	leaves distributed regularly along the stem
			4	shoot scarcely foliated
Bud bank: vertical layers	2442	6052	5	tufts and crowns, leaves concentrated as a rosette at the top of taller shoot or stem
			6	other
			1	no buds per shoot (not applicable)
			1.1	no buds per shoot, below soil surface, < -10 cm
			1.2	no buds per shoot, below soil surface, 0 < x < -10 cm
			1.3	no buds per shoot at soil surface
			1.4	no buds per shoot, above soil surface, 0 > x > 10 cm
			1.5	no buds per shoot, above soil surface, > 10 cm
			2	1–10 buds per shoot
			2.1	1–10 buds per shoot, below soil surface, < -10 cm
			2.2	1–10 buds per shoot, below soil surface, 0 < x < -10 cm
			2.3	1–10 buds per shoot at soil surface
			2.4	1–10 buds per shoot, above soil surface, 0 > x > 10 cm
			2.5	1–10 buds per shoot, above soil surface, > 10 cm
			3	> 10 buds per shoot
3.1	> 10 buds per shoot, below soil surface, < -10 cm			
3.2	> 10 buds per shoot, below soil surface, 0 < x < -10 cm			
3.3	> 10 buds per shoot at soil surface			
3.4	> 10 buds per shoot, above soil surface, 0 > x > 10 cm			
3.5	> 10 buds per shoot, above soil surface, > 10 cm			
Bud bank – seasonality	2468	6203	1	seasonal
			1.1	seasonal, above-ground
			1.2	seasonal, below-ground
			2	perennial
			2.1	perennial, above-ground
			2.2	perennial, below-ground
			3	seasonal & potential
			3.1	seasonal & potential, above-ground
			3.2	seasonal & potential, below-ground
			4	perennial & potential
Clonal growth organs	1958	5540	4.1	perennial & potential, above-ground
			4.2	perennial & potential, below-ground
Life span of a shoot	1737	4233	17	17 categories hierarchical classified according to their placement (above, at or below soil surface) and again subdivided to their origin (stem, root or leaf origin) (see Data Standards)
			1	monocyclic (1 year)
Persistence of connection between parent and offspring shoots	1834	4683	2	dicyclic or polycyclic (> 1 year)
			1	< 1 year
			2	1–2 years
			3	> 2 years

Table 2. Continued

Trait name in data standards	Number of species	Number of records	Catalogue Number	Category or unit(s) of measurement			
Number of offspring shoots per parent shoot per year	1740	4263	1	< 1 shoot/parent shoot/year			
			2	1 shoot/parent shoot/year			
			3	2–10 shoots/parent shoot/year			
			4	> 10 shoots/parent shoot/year			
Lateral spread	555	1089	1	< 0.01 m year ⁻¹			
			2	0.01–0.25 m year ⁻¹			
			3	0.25 m year ⁻¹			
			4	dispersable diaspores			
Seed number	1767	6165		number of seeds per ramet			
Seed crop frequency	196	201	1	more than once a year			
			2	once a year			
			3	once in 2 years			
			4	once in > 2 years			
			5	not applicable			
			6	unknown			
Seed shedding	1640	3331		month of the year (1–12)			
Seed weight	2025	7239		mg			
Seed size	2401	6578		length, width and height (mm)			
Seed shape	2401	6578		calculated from seed length, width and height (unitless, see Data Standards)			
Soil seed bank type	1479	44353		transient – short-term persistent – long-term persistent			
Seed bank longevity index	1479	44353		short-lived (0) – long-lived (1)			
Soil seed bank density	1479	44353		per m ²			
Diaspore type categories	2082	4162	1	vegetative dispersule			
			2	generative dispersule			
			2.1	one-seeded			
			2.2	multi-seeded			
			3	germinule			
			4	unknown			
			Morphology of dispersal unit	2082	4162	1	nutrient containing structures
						2	elaiosome
						3	aril
						4	pulp
						5	balloon structures
						5.1	open balloons
5.2	closed balloons						
8	flat appendages						
8.1	small flat appendages						
8.2	large flat appendages						
9	elongated appendages						
9.1	one short elongated appendage						
9.2	two or more short elongated appendages						
9.3	one long elongated appendage						
9.4	two or more long elongated appendages						
9.1–4	additional info: hooked structures						
10	no appendages						
10.1	seed with coarse surface, no appendages						
10.2	seed with smooth surface, no appendages						
11	other specialisations						
12	unknown						
Seed release height	2586	3921		m			
Terminal velocity seeds	1328	2592		m s ⁻¹			
Buoyancy	989	8081		number or % of floating seeds			
Epizoochory	192	559		number or % of attached seeds			
Endozoochory	149	179		number or % seeds that survived ingestion			
Dispersal data obtained from literature	2956	13920		14 dispersal type categories (see Data Standards), 32 dispersal vector categories (see Data Standards)			
Habitat characteristics	1401	1401		Categories referring to soil moisture, acidity, substrate, type, nutrient status. Water column acidity, alkalinity, and sediment redox potential for aquatic plants			



Fig. 1. Geographical range covered by the data base. Dark grey: core regions; light grey: overlap > 50% with the national floras.

seed release height is important in modelling wind dispersal. One key factor in epizoochory (dispersal by means of seeds attached to external parts of an animal) is the capacity of seeds to remain attached to fur, i.e. the attachment potential (Couvreur *et al.* 2004; Römermann *et al.* 2005). Other dispersal traits covered by the LEDA Traitbase include endozoochory (seeds dispersed after passing through the digestive tract of an animal), buoyancy (floating capacity), morphology of the dispersal unit and information about dispersal types as well as dispersal vectors of plants.

THE LEDA GEOGRAPHICAL RANGE AND TAXONOMIC CORE

The geographical range of the LEDA project (Fig. 1) roughly covers NW-Europe from the North Cape, Norway, to the Loire in France, and from the eastern borders of both Finland and Germany to the west coast of Ireland. Plant species present in Austria, Switzerland, Iceland, Poland, the Baltic States, Czech Republic, Slovakia and Hungary overlap with those in the core LEDA area by 50–80%, indicating the wide range of possible users of the Traitbase.

Selection of the 3000 priority vascular plant species for which we collected data was made according to the species frequencies in the core countries, i.e. UK, The Netherlands and Germany, disregarding alpine species and extremely rare species.

The taxonomic core of the LEDA Traitbase consists of one synonymised plant list at the species level, complete with authorities. The list was collated from the national plant lists available for the geographical range of the LEDA project (see Appendix S1). Species names and grouping of the species in higher taxa, however, cannot be considered as a stable

reference system because taxonomies are subject to research and are changed frequently. When collating existing data bases and retrieving data from published literature different 'taxonomic concepts' (*sensu* Geoffroy & Berendsohn 2003) inevitably get merged. The resulting loss in data quality can, however, be expected have little impact on the LEDA trait data base for the following reasons: (1) Only a few taxonomic groups in the flora of Northwest Europe are still under profound revision, and (2) Floras were used, which are interconnected by the species checklist from the SynBioSys-Europe project (Schaminée *et al.* 2007).

METHODS OF COLLECTING DATA

In LEDA, the following trait data bases were collated: Ecoflora (Fitter & Peat 1994), Electronic Comparative Plant Ecology (Hodgson *et al.* 1995), Biological traits of vascular plants data base (Kleyer 1995), CLOPLA (Klimešová & Klimeš 2006), the Soil seed bank data base (Thompson *et al.* 1997), the Dutch Botanical Database (CBS 1997 with updates), DIASPORUS (Bonn *et al.* 2000), seed mass data from BioFlor (Klotz *et al.* 2002), and BioPop (Poschlod *et al.* 2003; Jackel *et al.* 2006).

The remaining data were derived from literature dating back to the 19th Century. For many traits in the LEDA Traitbase, we expected more data to be available in the literature than we were actually able to retrieve. A large field sampling campaign was used to obtain data identified as missing in the literature: collecting and measuring standards are described in Knevel *et al.* (2005, www.leda-traitbase.org; see also Cornelissen *et al.* 2003). Very rare species had to be excluded because sampling effort increased with rarity or because extraction of plant material from the field was prohibited by conservation authorities. Age of first flowering could not be determined during field collections and was therefore compiled solely from the literature.

The LEDA editorial board ensures that each entry in the Traitbase has a full reference to its original source, whether a published book, article, data base or recent measurement according to the LEDA standards. Also, newly measured data are referenced according to the field site, including georeference information and habitat characteristics. When habitat characteristics were missing for data from other sources, these were derived from indicator values (Ellenberg *et al.* 1991).

TECHNICAL STRUCTURE OF THE DATA BASE

The LEDA Traitbase is a combination of a relational data base holding the trait data and several web applications allowing for input, access and analysis of trait-related data (see Appendix S1). Users only need a Web browser to query the LEDA Traitbase. After query execution, a table containing the selected records will be displayed within the web browser, either as individual records with bibliographic reference or as aggregated values, e.g. the average of all SLA records for a species. Registered users may upload or otherwise compose a

list of species names to constrain their queries, and they may instruct the system to deliver the query results to their e-mail address. For example, to obtain a GIS link, the mapping module FloraMap of the German online plant atlas (<http://www.floraweb.de>) is accessible from within the LEDA web query application. This allows access to further trait data, species distribution data and related information (in German for version 1). Hence, the analysis of the spatial distribution of traits (e.g. Kühn *et al.* 2006) can be facilitated easily.

Applications of the LEDA Traitbase

Potential applications of the LEDA Traitbase cover the whole range of functional ecology and phylogenetic ecology, the fusion of ecology and evolutionary history (e.g. Grime 2006; McGill *et al.* 2006; Westoby 2006). A major field of functional ecology is the analysis of changes in community trait composition in response to environmental change to reveal functional response traits (Lavorel & Garnier 2002). Understanding how persistence, regeneration and dispersal traits respond to environmental change is essential for the prediction of species change in many ecological applications (e.g. landscape planning, restoration, mitigation of plant invasions). For instance, the LEDA Traitbase has been used to show that the predictability of local species composition from environmental conditions is constrained by dispersal traits (Ozinga *et al.* 2005). Dispersal traits were further used to assess wind dispersal potential or external animal dispersal in plants (Tackenberg *et al.* 2003). LEDA Traitbase data were also used to model relationships between plant traits, soil fertility and disturbance by land use (Kleyer 2002; Kühner & Kleyer 2008), and bud bank traits were used to explain the regeneration of biennials and perennials following disturbance of urban plant communities (Latzel *et al.* 2008).

Potential applications of the LEDA Traitbase include the analysis of changes in ecosystem functions (e.g. productivity, carbon sequestration) in response to changes in biodiversity and community composition based on the concomitant changes in 'functional effect traits'. Changes in traits such as longevity, leaf dry matter content, leaf nitrogen content or woodiness can affect the productivity of plant communities (Garnier *et al.* 2007), nutrient cycling (Eviner *et al.* 2006), or soil carbon sequestration (De Deyn *et al.* 2008). Response and effect traits are linked when changes in species composition translate into modifications of ecosystem properties (Chapin *et al.* 2000). For instance, seed production may be essential for the response of plant species to strong disturbances and at the same time an essential resource for animals. On the other hand, while seeds may be important for the response to disturbance, leaf and stem traits may be more important for the effect of plant species on biomass decomposition. By coupling the LEDA Traitbase with data sets that combine species abundances with environmental information and ecosystem properties, response and effect traits and linkages between these can be identified (Suding *et al.* 2008).

LEDA data were also used for the analysis of relationships between traits and distribution patterns of rarity and endan-

germent of plant species (Smart *et al.* 2005; Römermann *et al.* 2008). Specifically, there has been a long quest for traits that make species invasive (e.g. Kühn *et al.* 2004; Moles *et al.* 2008, see Pyšek & Richardson 2007 for a review) or influence commonness and rarity in weeds (e.g. Lososová *et al.* 2008) or urban plant species (Thompson & McCarthy, in press).

Functional diversity, i.e. the value and range of plant functional traits in a given community (Tilman 2001), has been proposed as an important feature of communities, for instance to provide resilience in relation to regime shifts in terrestrial and aquatic communities (Folke *et al.* 2004). Functional diversity can be recorded at different biological levels, e.g. within species and between species in a community. Intraspecific diversity can be extracted from the LEDA Traitbase either by retrieving the original individual records or from aggregated information such as minimum and maximum values or standard deviations. For rare species or native species of natural landscapes without agricultural land use, the number of trait records is still small. More records will be needed throughout the geographical and environmental range of the species to assess the full extent of trait variability. Interspecific diversity can be measured with various indices (e.g. Mason *et al.* 2005) by collating LEDA data aggregated per trait and species to vegetation relevés.

Understanding how investments of carbon and mineral nutrients vary between species is central to plant ecology. Large data bases on plant traits have helped to clarify the extent to which scaling relations between traits indicate potential trade-offs or allometries (e.g. Enquist & Niklas 2002; Wright *et al.* 2004). Although LEDA comprises only limited data on biomass partitioning, it can produce trait correlation structures that could assist in revealing scaling relations associated with persistence, regeneration and dispersal. In contrast to such physiologically determined trade-offs, environment-induced trade-offs are often characterised by different costs and benefits along environmental gradients. LEDA data have been used to search for trade-offs between local above-ground persistence and below-ground seed persistence (Ozinga *et al.* 2007) and between generative and vegetative reproduction in riparian vegetation (Boedeltje *et al.* 2008).

These examples show that the LEDA Traitbase can assist in clarifying the role of traits and of trait variation in the response of plants to changing environments, the assembly of communities and the functioning of ecosystems. Case studies exploring these issues will most often take place at the level of a community or a landscape. Trait measurements at these levels will profit from assessing the variation of the traits under study against variation in the flora of the region or biome. This information can now be retrieved from the LEDA Traitbase for the flora of NW Europe. The LEDA Traitbase also offers the opportunity to re-analyse large vegetation data sets in terms of functional traits. For instance, it would be interesting to combine country-wide sets of relevés aggregated to syntaxonomic classes (e.g. Schaminée *et al.* 1995–1999) with the LEDA Traitbase to extract variation

in persistence and regeneration traits of plant communities. So far, this has only been done for dispersal traits (Ozinga *et al.* 2005). Such community trait profiles could be used to generate better hypotheses for detailed investigations of plant trait–environment linkages (McGill *et al.* 2006).

Further prospects

At present the Traitbase supports a total of more than 8300 taxa of NW-Europe. Many taxa are subspecies to which no data are linked. However, the possibility exists to link data to these taxa, as well as to taxa that currently are not included in the LEDA priority list. The LEDA consortium welcomes new collaborators interested in delivering new data to the LEDA Traitbase. The LEDA standards (available at through www.leda-traitbase.org) provide baseline information on how the data should be organised. To assure data quality and consistency with the LEDA data standard, the LEDA Editorial Board will review the data before incorporating them into LEDA.

The LEDA Traitbase and its applications are designed to be extended with further traits. Adapting the data base scheme is relatively easy, since data for distinct traits are stored within distinct tables. The LEDA consortium welcomes any initiative that seeks to enlarge the LEDA Traitbase, either by extension of the geographical range or by extension of the traits that are covered by the data base. This would include the obligation to establish appropriate data standards, support additional technical effort and to take part in the reviewing process.

Moreover, we see future prospects in the collation of LEDA to various other data bases, such as plant genomics, distribution, Red Lists, plant communities, habitats and environmental factors, e.g. nutrient and disturbance data for sites with known species composition (Bekker *et al.* 2007; Schaminée *et al.* 2007). Currently, there are many initiatives across Europe and other parts of the world that intend to make available various data bases. We expect that the joint analysis of data from these different sources will greatly advance our understanding of large-scale biodiversity change.

Acknowledgement

The authors thank Heike Zimmermann, Tobias Kerrinnes, Hartmut Fridl, Susanne Bonn, Anne Krämer, Inge Lauer, Stefan Reidinger, Carsten Rüter, Tomas Flower-Ellis, Franciska Schultz, Espen Aarnes, Christina Lüst, Melanie Siemon, Christoph Reiffert and Svenja Vozenilek who assisted in measurements or data input.

The LEDA Traitbase project was funded by the European Commission through the fifth framework under the EESD programme, project number EVR1-CT-2002-40022. The authors gratefully acknowledge the help of Hans Fink and Rudolf May, German Federal Agency for Nature Conservation, Bonn, DE, in hosting and maintaining LEDA Traitbase after the end of the funding period.

References

Barkman, J. (1988) New Systems of Plant growth forms and phenological plant types. *Plant Form and Vegetation Structure* (eds M.J.A. Werger, P.J.M. Van der Aart, H.J. Doring & J.T.A. Verhoeven), pp. 9–44. S. P. B. Publishers, The Hague.

- Bekker, R.M., Schaminée, J.H.J., Bakker, J.P. & Thompson, K. (1998) Seed bank characteristics of Dutch plant communities. *Acta Botanica Neerlandica*, **47**, 15–26.
- Bekker, R.M., van der Maarel, E., Bruelheide, H. & Woods, K. (2007) Long-term datasets: from descriptive to predictive data using ecoinformatics. *Journal of Vegetation Science*, **18**, 458–462.
- Bellingham, P.J. & Sparrow, A.D. (2000) Resprouting as a life history strategy in woody plant communities. *Oikos*, **89**, 409–416.
- Boedeltje, G., Ozinga, W.A., & Prinzing, A. (2008) The trade-off between vegetative and generative reproduction among angiosperms influences regional hydrochorous propagule pressure. *Global Ecology and Biogeography*, **17**, 50–58.
- Bonn, S., Poschlod, P. & Tackenberg, O. (2000) Diasporus – a database for diaspore dispersal – concept and application in case studies for risk assessment. *Zeitschrift für Ökologie und Naturschutz*, **9**, 85–97.
- Bruun, H.H. & Poschlod, P. (2006) Why are small seeds dispersed through animal guts: large numbers or seed size per se? *Oikos*, **113**, 403–411.
- Cain, M.L., Milligan, B.G. & Strand, A.E. (2000) Long-distance seed dispersal in plant populations. *American Journal of Botany*, **87**, 1217–1227.
- CBS (1997) *Dutch Botanical Database, Biobase*. Centraal Bureau voor Statistiek, Voorburg/Heerlen.
- Chapin, F.S., Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L. *et al.* (2000) Consequences of changing biodiversity. *Nature*, **405**, 234–242.
- Cornelissen, J.H.C., Lavorel, S., Garnier, E., Diaz, S., Buchmann, N., Gurvich, D.E. *et al.* (2003) Handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian Journal of Botany*, **51**, 335–380.
- Couvreur, M., Vandenberghe, B., Verheyen, K. & Hermy, M. (2004) Experimental assessment of seed adhesivity on animal furs. *Seed Science Research*, **14**, 147–159.
- Danvind, M. & Nilsson, C. (1997) Seed floating ability and distribution of alpine plants along a northern Swedish river. *Journal of Vegetation Science*, **8**, 271–276.
- De Deyn, G.B., Cornelissen, J.H.C. & Bardgett, R.D. (2008) Plant functional traits and soil carbon sequestration in contrasting biomes. *Ecology Letters*, **11**, 516–531.
- De Kroon, H. & van Groenendael, J. (eds) (1997) *The Ecology and Evolution of Clonal Plants*. Backhuys Publishers, Leiden.
- Du Rietz, G.E. (1931) Life forms of terrestrial flowering plants. *Acta Phytogeographica Suecica*, **3**, 1–95.
- Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W. & Paulissen, D. (1991) Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobotanica*, **18**, 1–248.
- Enquist, B.J. & Niklas, K.J. (2002) Global allocation rules for patterns of biomass partitioning in seed plants. *Science*, **295**, 1517–1520.
- Eviner, V.T., Chapin, F.S. & Vaughn, C.E. (2006) Seasonal variations in plant species effects on soil N and P dynamics. *Ecology*, **87**, 974–986.
- Fitter, A.H. & Peat, H.J. (1994) The ecological flora data-base. *Journal of Ecology*, **82**, 415–425.
- Flynn, S., Turner, R.M. & Stuppy, W.H. (2006) *Seed Information Database* (release 7.0, October 2006) Available: <http://www.kew.org/data/sid>.
- Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L. & Holling, C.S. (2004) Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology and Systematics*, **35**, 557–581.
- Garnier, E., Lavorel, S., Ansquer, P., Castro, H., Cruz, P., Dolezal, J. *et al.* (2007) Assessing the effects of land-use change on plant traits, communities and ecosystem functioning in grasslands: a standardized methodology and lessons from an application to 11 European sites. *Annals of Botany*, **99**, 967–985.
- Geoffroy, M. & Berendsohn, W.G. (2003) The concept problem in taxonomy: importance, components, approaches. *Schriftenreihe Vegetationsk.* **39**, 5–14.
- Grime, J.P. (2006) Trait convergence and trait divergence in herbaceous plant communities: mechanisms and consequences. *Journal of Vegetation Science*, **17**, 255–260.
- Grime, J.P., Hodgson, J.G. & Hunt, R. (1988) *Comparative Plant Ecology*. Unwin Hyman, London.
- Hodgson, J.G., Grime, J.P., Hunt, R. & Thompson, K. (1995) *The Electronic Comparative Plant Ecology*. Chapman & Hall, London.
- Jackel, A.-K., Dannemann, A., Tackenberg, O., Kleyer, M. & Poschlod, P. (2006) BioPop – Funktionelle Merkmale von Pflanzen und ihre Anwendungsmöglichkeiten im Arten-, Biotop- und Naturschutz. *Naturschutz und Biologische Vielfalt*, **32**, 1–168.
- Kirchner, O., Loew, E. & Schroeter, C. (1908–1936) *Lebensgeschichte der Blütenpflanzen Mitteleuropas*. Ulmer, Stuttgart.

- Kleyer, M. (1995) *Biological Traits of Vascular Plants – A Database*. University of Stuttgart, Arbeitsberichte, N.F. 2, Stuttgart, Germany.
- Kleyer, M. (2002) Validation of plant functional types across two contrasting landscapes. *Journal of Vegetation Science*, **13**, 167–178.
- Klímeš, L., Klímešová, J., Hendriks, R. & Van Groenendael, J.M. (1997) Clonal plant architectures: a comparative analysis of form and function. *The ecology and evolution of clonal plants* (eds H. de Kroon & J. van Groenendael), pp. 1–29. Backhuys Publishers, Leiden.
- Klímeš, L. & Klímešová, J. (2000) Plant rarity and the mode of clonal growth. *Zeitschrift für Ökologie und Naturschutz*, **9**, 43–52.
- Klímešová, J. & Klímeš, L. (2006) *CLO-PLA3*. <http://clopla.butbn.cas.cz/>.
- Klímešová, J. & Klímeš, L. (2007) Budbanks and their role in vegetation regeneration. A literature review and proposal for simple classification. *PPEES* **8**, 115–129.
- Klímešová, J. & Martinková, J. (2004) Intermediate growth forms as a model for the study of plant clonality functioning: an example with root sprouters. *Evolutionary Ecology*, **18**, 669–681.
- Klotz, S., Kühn, I. & Durka, W. (2002) BIOLFLOR – Eine Datenbank zu biologisch-ökologischen Merkmalen der Gefäßpflanzen in Deutschland. *Schriftenreihe für Vegetationskunde*, **38**. Bundesamt für Naturschutz, Bonn.
- Knevel, I.C., Bekker, R.M., Bakker, J.P. & Kleyer, M. (2003) Life-history traits of the NW European flora: the LEDA database. *Journal of Vegetation Science*, **14**, 611–614.
- Knevel, I.C., Bekker, R.M., Kunzmann, D., Stadler, M. & Thompson, K. (2005) *The LEDA Traitbase Collecting and Measuring Standards of Life-history Traits of the NW European Flora*. Published by the University of Groningen, Groningen, The Netherlands. Available: <http://www.LEDA-Traitbase.org>.
- Kühn, I., Bierman, S.M., Durka, W. & Klotz, S. (2006) Relating geographical variation in pollination types to environmental and spatial factors using novel statistical methods. *New Phytologist*, **172**, 127–139.
- Kühn, I., Brandenburg, M. & Klotz, S. (2004) Why do alien plant species that reproduce in natural habitats occur more frequently? *Diversity & Distributions*, **10**, 417–425.
- Kühner, A. & Kleyer, M. (2008) A parsimonious combination of traits predicting plant response to disturbance and soil fertility. *Journal of Vegetation Science*, **19**, 681–692.
- Latzel, V., Mihaluk, S., Klímešová, J. (2008) Plant traits and regeneration of urban plant community after disturbance: does bud bank play any role? *Journal of Applied Vegetation Sciences First Online* (doi: 10.3170/2008-7-18487).
- Lavorel, S. & Garnier, E. (2002) Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional Ecology*, **16**, 545–556.
- Leishman, M.R. (2001) Does the seed size/number trade-off model determine plant community structure? An assessment of the model mechanisms and their generality. *Oikos*, **93**, 294–302.
- Lososová, Z., Chytrý, M., Kühn, I. (2008) Plant attributes determining regional commonness and rarity of weeds on Central European arable land. *Journal of Biogeography*, **35**, 177–187.
- Mason, N.W.H., Mouillod, D., Lee, W.G. & Wilson, J.B. (2005) Functional richness, functional evenness and functional divergence: the primary components of functional diversity. *Oikos*, **111**, 112–118.
- McGill, B.J., Enquist, B.J., Weiher, E., & Westoby, M. (2006) Rebuilding community ecology from functional traits. *Trends in Ecology & Evolution*, **21**, 178–185.
- Moles, A.T., Gruber, M.A.M. & Bonser, S.P. (2008) A new framework for predicting invasive plant species. *Journal of Ecology*, **96**, 13–17.
- Nathan, R., Katul, G.G., Horn, H.S., Thomas, S.M., Oren, R., Avissar, R., Pacala, S.W. & Levin, S.A. (2002) Mechanisms of long-distance dispersal of seeds by wind. *Nature*, **418**, 409–413.
- Ozinga, W.A., Hennekens, S.M., Schaminée, J.H.J., Smits, N.A.C., Bekker, R.M., Romermann, C., Klimes, L., Bakker, J.P. & van Groenendael, J.M. (2007) Local above-ground persistence of vascular plants: life-history trade-offs and environmental constraints. *Journal of Vegetation Science*, **18**, 489–497.
- Ozinga, W.A., Schaminée, J.H.J., Bekker, R.M., Bonn, S., Poschlod, P., Tackenberg, O., Bakker, J.P. & van Groenendael, J.M. (2005) Predictability of plant species composition from environmental conditions is constrained by dispersal limitation. *Oikos*, **108**, 555–561.
- Poschlod, P., Kleyer, M., Jackel, A.-K., Dannemann, A. & Tackenberg, O. (2003) BIOPOP – a database of plant traits and internet application for nature conservation. *Folia Geobotanica*, **38**, 263–271.
- Poschlod, P., Tackenberg, O. & Bonn, S. (2005) Plant dispersal potential and its relation to species frequency and coexistence. (ed. E. van der Maarel), pp. 147–171. *Vegetation Ecology*. Blackwell, Oxford.
- Pyšek, P. & Richardson, D.M. (2007) Traits associated with invasiveness in alien plants: where do we stand? *Biological Invasions*, ed. W. Nentwig. *Ecological Studies*, **193**, 97–126. Springer, Berlin & Heidelberg.
- Rabotnov, T.A. (ed.) (1974–1990) *Biological Flora of the Moscow Region*, Volumes 1–8. Izdatelstvo Moskovskogo Universiteta, Moskva.
- Raunkiaer, C. (1937) *Plant Life Forms*. Clarendon Press, Oxford.
- Rejmánek, M. & Richardson, D.M. (1996) What attributes make some plant species more invasive? *Ecology*, **77**, 1655–1661.
- Römermann, C., Tackenberg, O., Jackel, A.-K. & Poschlod, P. (2008) Eutrophication and fragmentation are related to species' rate of decline but not to species rarity – results from a functional approach. *Biodiversity and Conservation*, **17**, 591–604.
- Römermann, C., Tackenberg, O. & Poschlod, P. (2005) How to predict attachment potential of seeds to sheep and cattle coat from simple morphological seed traits. *Oikos*, **110**, 219–230.
- Ryser, P. (1996) The importance of tissue density for growth and lifespan of leaves and roots: a comparison of five ecologically contrasting grasses. *Functional Ecology*, **10**, 717–723.
- Schaminée, J.H.J., Hennekens, S.M., & Ozinga, W.A. (2007) Use of the ecological information system SynBioSys for the analysis of large datasets. *Journal of Vegetation Science*, **18**, 463–470.
- Schaminée, J.H.J., Hommel, P.W.F.M., Stortelder, A.H.F., Weeda, E.J. & Westhoff, V. (1995–1999) *De Vegetatie van Nederland*, Vol. 1–5. Opulus Press, Uppsala/Leiden.
- Smart, S.M., Bunce, R.G.H., Marrs, R., LeDuc, M., Firbank, L.G., Maskell, L.C., Scott, W.A., Thompson, K. & Walker, K.J. (2005) Large-scale changes in the abundance of common higher plant species across Britain between 1978, 1990 and 1998 as a consequence of human activity: tests of hypothesised changes in trait representation. *Biological Conservation*, **124**, 355–371.
- Suding, K.N., Lavorel, S., Chapin, F.S., Cornelissen, J.H.C., Diaz, S., Garnier, E., Goldberg, D., Hooper, D.U., Jackson, S.T. & Navas, M.L. (2008) Scaling environmental change through the community-level: a trait-based response-and-effect framework for plants. *Global Change Biology*, **14**, 1125–1140.
- Tackenberg, O., Poschlod, P. & Bonn, S. (2003) Assessment of wind dispersal potential in plant species. *Ecological Monographs*, **73**, 191–205.
- Thompson, K., Bakker, J.P. & Bekker, R.M. (1997) *Soil Seed Banks of North West Europe: Methodology, Density and Longevity*. Cambridge University Press, Cambridge.
- Thompson, K. & McCarthy, M.A. (2005) Traits of British alien and native urban plants. *Journal of Ecology*, **96**, 853–859.
- Tilman, D. (2001) Functional diversity. *Encyclopedia of biodiversity*. (ed. S.A. Levin), pp. 109–120. Academic Press, San Diego, CA.
- USDA, NRCS. (2006) *The PLANTS Database* (<http://plants.usda.gov>, 7 November 2006). National Plant Data Center, Baton Rouge, LA.
- Van der Pijl, L. (1972) *Principles of Dispersal in Higher Plants*. Springer, Berlin.
- Vesk, P.A. & Westoby, M. (2004) Sprouting by plants: the effect of modular organization. *Functional Ecology*, **18**, 939–946.
- Westoby, M. (2006) Phylogenetic ecology at world scale, a new fusion between ecology and evolution. *Ecology*, **87**, S163–S166.
- Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. (2002) Plant ecological strategies: some leading dimensions of variation between species. *Annual Review of Ecology and Systematics*, **33**, 125–159.
- Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F. et al. (2004) The worldwide leaf economics spectrum. *Nature*, **428**, 821–827.

Received 3 December 2007; accepted 3 July 2008

Handling Editor: Michael Hutchings

Supporting information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Details regarding the LEDA trait definitions and the structure of the data base.

Please note: Blackwell Publishing are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.